

The Infectious Cycle

Lecture 2

Biology 3310/4310


Virology

Spring 2017

"You know my methods, Watson"
--SIR ARTHUR CONAN DOYLE


The Infectious Cycle

Virologists divide the infectious cycle into steps to facilitate their study, but no such artificial boundaries occur




Some important definitions

- A **susceptible** cell has a functional receptor for a given virus - *the cell may or may not be able to support viral replication*
- A **resistant** cell has no receptor - *it may or may not be competent to support viral replication*
- A **permissive** cell has the capacity to replicate virus - *it may or may not be susceptible*
- A **susceptible AND permissive** cell is the only cell that can take up a virus particle and replicate it



- Animal viruses at first could not be routinely propagated in cultured cells
- Most viruses were grown in laboratory animals






Embryonated eggs at 10 to 12 days being inoculated by automated machinery. 1st larger needle (about 1 mm diameter) punches a hole in a shell and 2nd smaller needle injects a seed into the allantoic cavity of the egg followed by incubation for 2 to 3 days. It takes less than 10 seconds to inoculate a row of eggs.
Courtesy: Solvay




Studying the infectious cycle in cells


- Not possible before 1949 (animal viruses)
- *Enders, Weller, Robbins* propagate poliovirus in human cell culture - primary cultures of embryonic tissues
- Nobel prize, 1954




Virus cultivation



Primary human
foreskin fibroblasts



Mouse fibroblast
cell line (3T3)




Human epithelial
cell line (HeLa)

continuous cell lines

Diploid cell strains (e.g. WI-38, human embryonic lung)











Go to:

m.socrative.com
room number: virus



A _____ and _____ cell is the only cell that can take up a virus particle and replicate it (fill in the blanks)

- A. Naive and resistant
- B. Primary and permissive
- C. Susceptible and permissive
- D. Susceptible and naive
- E. Continuous and immortal



cytopathic effect (CPE)

Formation of syncytia



Examples of cytopathic effects


Cytopathic effect(s)	Virus(es)
Morphological alterations	
Nuclear shrinking (pyknosis), proliferation of membrane	Picornaviruses
Proliferation of nuclear membrane	Alphaviruses, herpesviruses
Vacuoles in cytoplasm	Polyomaviruses, papillomaviruses
Syncytium formation (cell fusion)	Paramyxoviruses, coronaviruses
Margination and breaking of chromosomes	Herpesviruses
Rounding up and detachment of cultured cells	Herpesviruses, rhabdoviruses, adenoviruses, picornaviruses
Inclusion bodies	
Virions in nucleus	Adenoviruses
Virions in cytoplasm (Negri bodies)	Rabies virus
“Factories” in cytoplasm (Guarnieri bodies)	Poxviruses
Clumps of ribosomes in virions	Arenaviruses
Clumps of chromatin in nucleus	Herpesviruses


How many viruses in a sample?

- Infectivity
- Physical: virus particles and their components


Plaque assay

1930s: used to study multiplication of bacteriophages





Plaque assay



*1952, Renato Dulbecco developed for animal viruses
Nobel Prize, 1975*

PRODUCTION OF PLAQUES IN MONOLAYER TISSUE CULTURES BY SINGLE PARTICLES OF AN ANIMAL VIRUS

BY RENATO DULBECCO


CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CALIFORNIA




Read before the Academy, April 29, 1952


Research on the growth characteristics and genetic properties of animal viruses has stood greatly in need of improved quantitative techniques, such as those used in the related field of bacteriophage studies.

The requirements for a quantitative virus technique are as follows: (1) The use of a uniform type of host cell; (2) an accurate assay technique; (3) the isolation of the progeny of a single virus particle; and (4) the separate isolation of each of the virus particles produced by a single infected

Plaque assay







virology.ws/2013/06/15/the-wall-of-polio/




Go to:


m.socrative.com
room number: virus

When doing a plaque assay, what is the purpose of adding a semi-solid agar overlay on the monolayer of infected cells?


- A. To stabilize progeny virions
- B. To ensure that cells remain susceptible and permissive
- C. To act as a pH indicator
- D. To keep cells adherent to the plate during incubation
- E. To restrict viral diffusion after lysis of infected cells



**How many viruses are needed
to form a plaque?**




Plaque purification



A method for producing clonal virus stocks
Usually done 3 times

Endpoint dilution assay



Virus dilution		Cytopathic effect									
10^{-2}	+	+	+	+	+	+	+	+	+	+	+
10^{-3}	+	+	+	+	+	+	+	+	+	+	+
10^{-4}	+	+	-	+	+	+	+	+	+	+	+
10^{-5}	-	+	+	-	+	-	-	+	-	-	+
10^{-6}	-	-	-	-	-	-	+	-	-	-	-
10^{-7}	-	-	-	-	-	-	-	-	-	-	-

Particle-to-PFU ratio

- # of *physical* particles ÷ # of *infectious* particles
- A single particle *can* initiate infection
- Not all viruses are successful
 - Damaged particles
 - Mutations
 - Complexity of infectious cycle
- Complicates study

Particle-to-PFU ratios of some animal viruses

Virus	Particle/PFU ratio
<i>Papillomaviridae</i>	
Papillomavirus	10,000
<i>Picornaviridae</i>	
Poliovirus	30–1,000
<i>Herpesviridae</i>	
Herpes simplex virus	50–200
<i>Polyomaviridae</i>	
Polyomavirus	38–50
Simian virus 40	100–200
<i>Adenoviridae</i>	20–100
<i>Poxviridae</i>	1–100
<i>Orthomyxoviridae</i>	
Influenza virus	20–50
<i>Reoviridae</i>	
Reovirus	10
<i>Alphaviridae</i>	
Semliki Forest virus	1–2

Go to:


m.socrative.com
room number: virus

In the 'particle to pfu ratio', 'particle' can best be described as:



- A. One of the proteins which makes up the virion
- B. A virus which may or may not be infectious
- C. A virus which is infectious
- D. A virus which is not infectious
- E. Elementary or composite

One-step growth cycle


- Ellis & Delbruck, 1939, studies on *E. coli* bacteriophages
- Adsorb
- Dilute culture
- Sample
- Assay




Single and multi-step growth cycles




Adenovirus type 5





Synchronous infection - key to one-step growth cycle

To achieve this, we need to infect all the cells - but how do we know?



Multiplicity of infection (MOI)

- Number of infectious particles ADDED per cell
- Not the number of infectious particles each cell *receives*
- Add 10^7 virus particles to 10^6 cells - MOI of 10 - each cell does NOT receive 10 virions

MOI

- Infection depends on the random collision of virions and cells
- When susceptible cells are mixed with virus, some cells are uninfected, some receive one, two, three or more particles
- The distribution of virus particles per cell is best described by the *Poisson distribution*

$$P(k) = e^{-m} m^k / k!$$

$P(k)$: fraction of cells infected by k virus particles

m : multiplicity of infection (moi)

uninfected cells: $P(0) = e^{-m}$

cells receiving 1 particle: $P(1) = m e^{-m}$

cells multiply infected: $P(>1) = 1 - e^{-m}(m+1)$

[obtained by subtracting from 1

{the sum of all probabilities for any value of k }

the probabilities $P(0)$ and $P(1)$]

Examples:

If 10^6 cells are infected at moi of 10:

45 cells are uninfected

450 cells receive 1 particle

the rest receive >1 particle

If 10^6 cells are infected at moi of 1:

37% of the cells are uninfected

37% of the cells receive 1 particle

26% receive >1 particle

If 10^6 cells are infected at moi of .001:

99.9% of the cells are uninfected

00.099% of the cells receive 1 particle (990)

00.0001% receive >1 particle

Go to:

m.socrative.com
room number: virus

If cells are infected at an MOI=10 in a one-step growth cycle experiment, in the growth curve you will likely see...


- A. Multiple bursts of virus release
- B. Multiple eclipse periods
- C. A single burst of virus release
- D. No burst of virus release
- E. Asynchronous infection

Physical measurements of virus particles


- Hemagglutination




- Electron microscopy




- Viral enzymes





- Serology





- Nucleic acids




Hemagglutination






Measurement of viral enzyme activity








Enzyme-linked immunosorbent assay (ELISA): detecting viral antigens or antibodies










Green fluorescent protein




A**B**

Polymerase chain reaction (PCR)



- Research
- Industry
- Diagnosis



Deep, high-throughput sequencing

- Metagenomics
- Identification of new viruses in environmental samples
- Identification of new pathogens
- Human genome: 10 yr/\$3B vs 1 day/\$1500



(this is not DHTS)

TWiV 196: An arena for snakes

AUGUST 19, 2012



Hosts: Vincent Racaniello, Alan Dove, Rich Condit, Dickson Despommier, Kathy Spindler, Mark Stenglein, and Joseph DeRisi

The TWiVites meet with Mark Stenglein and Joseph DeRisi to discuss their discovery of a novel arenavirus in snakes with inclusion body disease.

<http://www.microbe.tv/twiv/twiv-196-an-arena-for-snakes/>

TWiV 199: Of mice, ticks, and pigs

SEPTEMBER 16, 2012



Hosts: Vincent Racaniello, Alan Dove, Rich Condit, and Kathy Spindler

Vincent, Alan, Rich, and Kathy discuss recent outbreaks of hantavirus pulmonary syndrome in Yosemite National Park and novel swine-origin influenza in the US midwest, and isolation of the Heartland virus from two patients in Missouri with severe febrile illness.



Click the arrow above to play, or right-click

<http://www.microbe.tv/twiv-199-of-mice-ticks-and-pigs/>

Viruses and viral sequences

RESEARCH ARTICLE



Zoonotic Viruses Associated with Illegally Imported Wildlife Products



January 13, 2012, 8:10 AM

From the Jungle to J.F.K., Viruses Cross Borders in Monkey Meat

By RACHEL NUWER



Carlos Rodríguez/Wildlife Conservation Society

A set of Cercopithecus monkey limbs recently confiscated at O'Hare Airport in Chicago.



It's a familiar story: deep within the jungle, an intrepid explorer or hunter awakens something he shouldn't have. Maybe he bagged the wrong bat or came into contact with an ailing chimp. Whatever the origin, he unsuspectingly becomes host to something deadly and unseen and unwittingly carries it back home across the river or the ocean.

And then the pandemic begins.

<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0029505>